

[SLAC](#) | [Library](#) | [Other Databases](#) | [Conferences](#) | [Institutions](#) | [Experiments](#)

[Questions/Comments](#)



[Help](#)  
[Search HEP](#)

Database: **HEP (USPIRES-SLAC)**

Modify your search. Press ENTER to run search again.

Searching for an author? Try [HEPNAMES](#).

Like to see another format? [cites](#) [citesummary](#) [LaTeX](#)

Result: **190** documents found:

1) [Comparison of three-jet events in p anti-p collisions at  \$s^{1/2} = 1.8\text{-TeV}\$  to predictions from a next-to-leading order QCD calculation.](#)

By D. Acosta, et al.,

[hep-ex/0410018] (Oct 2004) 28p.

2) [Measurement of partial widths and search for direct cp violation.](#)

By CDF Collaboration

FERMILAB-PUB-04-148-E (Aug 2004) 7p.

3) [Measurement of W gamma and Z gamma production in p anti-p collisions at  \$s^{1/2} = 1.96\text{-TeV}\$ .](#)

By CDF II Collaboration

[hep-ex/0410008] FERMILAB-PUB-04-246-E (Oct 2004) 7p.

4) [Measurement of the t anti-t production cross section in p anti-p collisions at  \$s^{1/2} = 1.96\text{-TeV}\$  using kinematic fitting of b-tagged lepton + jet events.](#)

By CDF-II Collaboration

[hep-ex/0409029] FERMILAB-PUB-04-207-E (Sep 2004) 20p.

5) [First measurements of inclusive W and Z cross sections from run II of the Tevatron collider.](#)

By CDF II Collaboration

[hep-ex/0406078] FERMILAB-PUB-04-159-E (Jun 2004) 8p.

6) [Search for doubly-charged Higgs bosons decaying to dileptons in p anti-p collisions at  \$s^{1/2} = 1.96\text{-TeV}\$ .](#)

By CDF Collaboration

[hep-ex/0406073] FERMILAB-PUB-04-112-E (Jun 2004) 15p.

7) [Inclusive search for anomalous production of high-p\(T\) like-sign lepton pairs in p anti-p collisions at  \$s^{1/2} = 1.8\text{-TeV}\$ .](#)

By CDF Collaboration

Phys.Rev.Lett.93:061802,2004. [hep-ex/0405063]

8) [Measurement of the t anti-t production cross section in p anti-p collisions at  \$s^{1/2} = 1.96\text{-TeV}\$  using dilepton events.](#)

By CDF Collaboration

[hep-ex/0404036] FERMILAB-PUB-04-051-E (Apr 2004) 18p.

9) [Direct photon cross section with conversions at CDF.](#)

By CDF Collaboration

[hep-ex/0404022] FERMILAB-PUB-04-044-E (Apr 2004) 11p.

10) [The Underlying event in hard interactions at the Tevatron anti-p p collider.](#)

By CDF Collaboration

[hep-ex/0404004] FERMILAB-PUB-04-041-E (Apr 2004) 22p.

11) [Optimized search for single top quark production at the Fermilab tevatron.](#)

By CDF Collaboration

Phys.Rev.D69:052003,2004.

12) [Search for  \$B\_0\(s\) \rightarrow \mu^+ \mu^-\$  and  \$B\_0\(d\) \rightarrow \mu^+ \mu^-\$  decays in p anti-p collisions at  \$s^{1/2} = 1.96\text{-TeV}\$ .](#)

By CDF Collaboration

Phys.Rev.Lett.93:032001,2004. [hep-ex/0403032]

- 13) [Observation of the narrow state X\(3872\) ---> J / psi pi+ pi- in anti-p p collisions at s\\*\\*\(1/2\) = 1.96-TeV.](#)  
By CDF II Collaboration  
Phys.Rev.Lett.93:072001,2004, Phys.Rev.Lett.93:072001,2004. [hep-ex/0312021]
- 14) [Heavy flavor properties of jets produced in p anti-p interactions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.D69:072004,2004. [hep-ex/0311051]
- 15) [Measurement of the polar angle distribution of leptons from W boson decay as a function of the W transverse momentum in proton anti-proton collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.D70:032004,2004. [hep-ex/0311050]
- 16) [Combination of CDF and D0 results on W boson mass and width.](#)  
By CDF Collaboration  
[hep-ex/0311039] (Nov 2003) 26p.
- 17) [Inclusive double pomeron exchange at the Fermilab Tevatron anti-p p collider.](#)  
By CDF Collaboration  
[hep-ex/0311023] FERMILAB-PUB-03-043-E (Nov 2003) 7p.
- 18) [Measurement of the average time integrated mixing probability of b flavored hadrons produced at the Tevatron.](#)  
By CDF Collaboration  
Phys.Rev.D69:012002,2004. [hep-ex/0309030]
- 19) [Search for the flavor changing neutral current decay D0 ---> mu+ mu- in p anti-p collisions at s\\*\\*\(1/2\) = 1.96-TeV.](#)  
By CDF Collaboration  
Phys.Rev.D68:091101,2003. [hep-ex/0308059]
- 20) [Measurement of the mass difference m\(D\(s\)+\) m\(D+\) at CDF2.](#)  
By CDF II Collaboration  
Phys.Rev.D68:072004,2003. [hep-ex/0310043]
- 21) [Measurement of prompt charm meson production cross-sections in p anti-p collisions at s\\*\\*\(1/2\) = 1.96-TeV.](#)  
By CDF Collaboration  
Phys.Rev.Lett.91:241804,2003. [hep-ex/0307080]
- 22) [Search for associated production of Upsilon and vector boson in p anti-p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.Lett.90:221803,2003.
- 23) [Search for new particles at CDF II.](#)  
By On behalf of the CDF Collaboration  
[hep-ex/0305027] FERMILAB-CONF-03-170-E (May 2003) 7p.
- 24) [Search for pair production of scalar top quarks in R parity violating decay modes in p anti-p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.Lett.92:051803,2004. [hep-ex/0305010]
- 25) [Central pseudorapidity gaps in events with a leading anti-proton at the Fermilab Tevatron p anti-p collider.](#)  
By CDF Collaboration  
Phys.Rev.Lett.91:011802,2003. [hep-ex/0303011]
- 26) [Search for the supersymmetric partner of the top quark in dilepton events from p pbar collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.Lett.90:251801,2003. [hep-ex/0302009]
- 27) [Cross-section for forward J / psi production in p anti-p collisions at S = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.D66:092001,2002.
- 28) [Search for long lived charged massive particles in anti-p p collisions at s\\*\\*1/2 = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.Lett.90:131801,2003. [hep-ex/0211064]

29) [Physics at large  \$p\(T\)^{\*\*2}\$  and  \$Q^{\*\*2}\$ : Summary.](#)

By G. Moortgat-Pick, S. Rolli, A.F. Zarnecki.

Acta Phys.Polon.B33:3955-3981,2002. [hep-ph/0210368]

30) [Search for a W-prime boson decaying to a top and bottom quark pair in 1.8-TeV p anti-p collisions.](#)

By CDF Collaboration

Phys.Rev.Lett.90:081802,2003. [hep-ex/0209030]

31) [Fragmentation Phenomena on QCD: Fragmentation Functions Approach and Parton Shower.](#)

By Simona Rolli.

FERMILAB-THESIS-1996-58 (Jan 1996) 166p.

32) [Search for radiative b hadron decays in proton anti-proton collisions at  \$s^{\*\*}\(1/2\) = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.D66:112002,2002. [hep-ex/0208035]

33) [Status of the CDF II experiment.](#)

By Simona Rolli.

FERMILAB-CONF-02-154-E (Aug 2002) 11p.

34) [Momentum distribution of charged particles in jets in dijet events in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\$ -TeV and comparisons to perturbative QCD predictions.](#)

By CDF Collaboration

Phys.Rev.D68:012003,2003.

35) [Measurement of the ratio of b quark production cross-sections in anti-p p collisions at  \$s^{\*\*}\(1/2\) = 630\$ -GeV and  \$s^{\*\*}\(1/2\) = 1800\$ -GeV.](#)

By CDF Collaboration

Phys.Rev.D66:032002,2002. [hep-ex/0206019]

36) [Branching ratio measurements of exclusive B+ decays to charmonium with the Collider Detector at Fermilab.](#)

By CDF Collaboration

Phys.Rev.D66:052005,2002.

37) [Limits on extra dimensions and new particle production in the exclusive photon and missing energy signature in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.89:281801,2002. [hep-ex/0205057]

38) [Measurement of B meson lifetimes using fully reconstructed B decays produced in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.D65:092009,2002.

39) [Search for new physics in photon lepton events in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.89:041802,2002. [hep-ex/0202044]

40) [Upsilon production and polarization in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.88:161802,2002.

41) [Comparison of the isolated direct photon cross-sections in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\$ -TeV and  \$s^{\*\*}\(1/2\) = 0.63\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.D65:112003,2002. [hep-ex/0201004]

42) [Distributed applications monitoring at system and network level.](#)

By MONARC Collaboration

Comput.Phys.Commun.140:219-225,2001.

43) [Soft and hard interactions in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1800\$ -GeV and 630-GeV.](#)

By CDF Collaboration

Phys.Rev.D65:072005,2002.

44) [Measurement of the B+ total cross-section and B+ differential cross-section  \$d\sigma / dp\(T\)\$  in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.D65:052005,2002. [hep-ph/0111359]

- 45) [Search for single top quark production in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D65:091102,2002. [hep-ex/0110067]
- 46) [Search for the decay  \$B/s \rightarrow \mu^+ \mu^- \Phi\$  in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D65:111101,2002.
- 47) [Charged jet evolution and the underlying event in proton-antiproton collisions at 1.8 TeV.](#)  
By CDF Collaboration  
Phys.Rev.D65:092002,2002.
- 48) [Search for new physics in photon lepton events in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D66:012004,2002. [hep-ex/0110015]
- 49) [Diffractive dijet production at  \$s^{\*\*}\(1/2\) = 630\text{-GeV}\$  and  \$1800\text{-GeV}\$  at the Fermilab Tevatron.](#)  
By CDF Collaboration  
Phys.Rev.Lett.88:151802,2002. [hep-ex/0109025]
- 50) [Study of the heavy flavor content of jets produced in association with W bosons in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D65:052007,2002. [hep-ex/0109012]
- 51) [Measurement of the strong coupling constant from inclusive jet production at the Tevatron anti-p p collider.](#)  
By CDF Collaboration  
Phys.Rev.Lett.88:042001,2002. [hep-ex/0108034]
- 52) [A Study of  \$B\_0 \rightarrow J/\psi K^0 \pi^+ \pi^-\$  decays with the collider detector at Fermilab.](#)  
By CDF Collaboration  
Phys.Rev.Lett.88:071801,2002. [hep-ex/0108022]
- 53) [Search for new heavy particles in the W Z0 final state in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.88:071806,2002. [hep-ex/0108004]
- 54) [Observation of diffractive J /  \$\psi\$  production at the Fermilab Tevatron.](#)  
By CDF Collaboration  
Phys.Rev.Lett.87:241802,2001. [hep-ex/0107071]
- 55) [Charged particle multiplicity in jets in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.87:211804,2001.
- 56) [Search for quark lepton compositeness and a heavy W-prime boson using the electron neutrino channel in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.87:231803,2001. [hep-ex/0107008]
- 57) [Searches for new physics in events with a photon and b-quark jet at CDF.](#)  
By CDF Collaboration  
Phys.Rev.D65:052006,2002. [hep-ex/0106012]
- 58) [Search for gluinos and squarks using like sign dileptons in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.87:251803,2001. [hep-ex/0106061]
- 59) [Measurement of  \$d\sigma/dM\$  forward backward charge asymmetry for high mass Drell-Yan  \$e^+ e^-\$  pairs from p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.87:131802,2001. [hep-ex/0106047]
- 60) [Cross-section and heavy quark composition of gamma + muon events produced in p anti-p collisions.](#)

By CDF Collaboration  
Phys.Rev.D65:012003,2002. [hep-ex/0106004]

61) [Search for gluinos and scalar quarks in p anti-p collisions at sqrt\(s\)=1.8 TeV using the missing energy plus multijets signature.](#)  
By CDF Collaboration  
Phys.Rev.Lett.88:041801,2002. [hep-ex/0106001]

62) [Production of chi/c1 and chi/c2 in p anti-p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.Lett.86:3963-3968,2001.

63) [Double diffraction dissociation at the Fermilab Tevatron collider.](#)  
By CDF Collaboration  
Phys.Rev.Lett.87:141802,2001. [hep-ex/0107070]

64) [Search for narrow diphoton resonances and for gamma gamma + w /z signatures in p anti p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.D64:092002,2001. [hep-ex/0105066]

65) [Measurement of the inclusive jet cross-section in anti-p p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.D64:032001,2001,Erratum-ibid.D65: 039903,2002. [hep-ph/0102074]

66) [Measurement of the t anti-t production cross-section in p anti-p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.D64:032002,2001,Erratum-ibid.D67: 119901,2003. [hep-ex/0101036]

67) [First measurement of the ratio B\(t ---> W b\) / B\(t ---> W q\) and associated limit on the CKM element |V\(tb\)|.](#)  
By CDF Collaboration  
Phys.Rev.Lett.86:3233-3238,2001. [hep-ex/0012029]

68) [The CDF Run II event data model.](#)  
By CDF Collaboration  
In \*CHEP 2000, Computing in high energy and nuclear physics\* 442-44.

69) [Atlas event data model optimization studies based on the use of segmented VArray in Objectivity/DB.](#)  
By S. Rolli, A. Salvadori, A.C. Schaffer, M. Schaller.  
In \*CHEP 2000, Computing in high energy and nuclear physics\* 436-44.

70) [Hardware trigger simulation at CDF.](#)  
By S. Rolli, J.D. Lewis, H. Ray, J. Nachtman, M. Worcester.  
In \*CHEP 2000, Computing in high energy and nuclear physics\* 250-25.

71) [Measurement of the two jet differential cross-section in proton anti-proton collisions at s\\*\\*\(1/2\) = 1800-GeV.](#)  
By CDF Collaboration  
Phys.Rev.D64:012001,2001,Erratum-ibid.D65: 039902,2002. [hep-ex/0012013]

72) [Measurement of the top quark p\(T\) distribution.](#)  
By CDF Collaboration  
Phys.Rev.Lett.87:102001,2001.

73) [Search for the supersymmetric partner of the top quark in p anti-p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.D63:091101,2001. [hep-ex/0011004]

74) [Search for neutral supersymmetric Higgs bosons in p anti-p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)  
By CDF Collaboration  
Phys.Rev.Lett.86:4472-4478,2001. [hep-ex/0010052]

75) [Test of enhanced leading order QCD in W boson plus jets events from 1.8-TeV anti-p p collisions.](#)  
By CDF Collaboration  
Phys.Rev.D63:072003,2001.

76) [Models of networked analysis at regional centres for LHC experiments \(MONARC\). Phase 2 report.](#)

By M. Aderholz, et al.,  
KEK-PREPRINT-2000-8 (Apr 2000) 43p.

77) [Measurement of the W boson mass with the collider detector at Fermilab.](#)

By CDF Collaboration  
Phys.Rev.D64:052001,2001. [hep-ex/0007044]

78) [Measurement of the decay amplitudes of  \$B^0 \rightarrow J/\psi K^\*\$  and  \$B^0\(S\) \rightarrow J/\psi \phi\$  decays.](#)

By CDF Collaboration  
Phys.Rev.Lett.85:4668-4673,2000. [hep-ex/0007034]

79) [Measurement of the top quark mass with the Collider Detector at Fermilab.](#)

By CDF Collaboration  
Phys.Rev.D63:032003,2001. [hep-ex/0006028]

80) [Measurement of  \$d\sigma/dy\$  for high mass Drell-Yan  \$e^+e^-\$  pairs from p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.D63:011101,2001. [hep-ex/0006025]

81) [Dijet production by double pomeron exchange at the Fermilab Tevatron.](#)

By CDF Collaboration  
Phys.Rev.Lett.85:4215-4220,2000.

82) [Measurement of J / psi and psi\(2S\) polarization in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.85:2886-2891,2000. [hep-ex/0004027]

83) [Search for color singlet technicolor particles in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.84:1110-1115,2000.

84) [Direct measurement of the W boson width in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.85:3347-3352,2000. [hep-ex/0004017]

85) [Search for second and third generation leptoquarks including production via technicolor interactions in p anti-p collisions at  \$\sqrt{s} = 1.8\$  TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.85:2056-2061,2000. [hep-ex/0004003]

86) [Limits on gravitino production and new processes with large missing transverse energy in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.85:1378-1383,2000. [hep-ex/0003026]

87) [Diffractive dijets with a leading anti-proton in anti-p p collisions at  \$\sqrt{s} = 1800\$ -GeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.84:5043-5048,2000.

88) [Search for new particles decaying to top - anti-top in proton - anti-proton collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.85:2062-2067,2000. [hep-ex/0003005]

89) [A Measurement of the differential dijet mass cross-section in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.D61:091101,2000. [hep-ex/9912022]

90) [Search for scalar top quark production in anti-p p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.84:5273-5278,2000. [hep-ex/9912018]

91) [Observation of orbitally excited B mesons in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.D64:072002,2001.

- 92) [Search for the charged Higgs boson in the decays of top quark pairs in the e tau and muon tau channels at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D62:012004,2000. [hep-ex/9912013]
- 93) [Search for scalar top and scalar bottom quarks in p anti-p collisions at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.84:5704-5709,2000. [hep-ex/9910049]
- 94) [Production of upsilon\(1S\) mesons from chi\(b\) decays in p anti-p collisions at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.84:2094-2099,2000. [hep-ex/9910025]
- 95) [Measurement of b quark fragmentation fractions in the production of strange and light B mesons in p anti-p collisions at  \$S^{\*\*}\(1/2\)=1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D60:092005,1999.
- 96) [Search for a W-prime boson via the decay mode W-prime ---> muon muon-neutrino in 1.8-TeV p - anti-p collisions.](#)  
By CDF Collaboration  
Phys.Rev.Lett.84:5716-5721,2000. [hep-ex/9910004]
- 97) [Measurement of the helicity of W bosons in top quark decays.](#)  
By CDF Collaboration  
Phys.Rev.Lett.84:216-221,2000. [hep-ex/9909042]
- 98) [Search for a fourth generation quark more massive than the Z0 boson in p anti-p collisions at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.84:835-840,2000. [hep-ex/9909027]
- 99) [Measurement of b quark fragmentation fractions in p anti-p collisions at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.84:1663-1668,2000. [hep-ex/9909011]
- 100) [A Measurement of sin 2 Beta from B ---> J / psi K0\(S\) with the CDF detector.](#)  
By CDF Collaboration  
Phys.Rev.D61:072005,2000. [hep-ex/9909003]
- 101) [The Transverse momentum and total cross-section of e+ e- pairs in the Z-boson region from p anti-p collisions at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.84:845-850,2000. [hep-ex/0001021]
- 102) [Observation of diffractive beauty production at the Fermilab Tevatron.](#)  
By CDF Collaboration  
Phys.Rev.Lett.84:232-237,2000.
- 103) [Measurement of the B0 anti-B0 oscillation frequency using l- D\\*+ pairs and lepton flavor tags.](#)  
By CDF Collaboration  
Phys.Rev.D60:112004,1999. [hep-ex/9907053]
- 104) [Searches for new phenomena at the Tevatron: SUSY and technicolor.](#)  
By CDF Collaboration  
In \*La Thuile 1999, Results and perspectives in particle physics\* 295-30.
- 105) [Search for technicolor particles in lepton plus two jets and multijet channels in p anti-p collisions at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
FERMILAB-PUB-99-141-E (May 1999) 10p.
- 106) [Search for the flavor changing neutral current decays B+ ---> muon+ muon- K+ and B0 ---> muon+ muon- K\\*0.](#)  
By CDF Collaboration  
Phys.Rev.Lett.83:3378-3383,1999. [hep-ex/9905004]
- 107) [Measurement of the B0\(D\) anti-B0\(D\) oscillation frequency using dimuon data in p anti-p collisions at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)  
By The CDF Collaboration  
Phys.Rev.D60:051101,1999.



108) [Light meson fragmentation functions.](#)

By S. Rolli.

In \*Minneapolis 1996, Particles and fields, vol. 1\* 599-60.

109) [Measurement of  \$B^0\$  - anti- \$B^0\$  flavor oscillations using jet charge and lepton flavor tagging in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.D60:072003,1999. [hep-ex/9903011]

110) [A Search for  \$B^0\(S\)\$  - anti- \$B^0\(S\)\$  oscillations using the semileptonic decay  \$B^0\(S\) \rightarrow \phi \text{ lepton}^+ X \text{ neutrino}\$ .](#)

By CDF Collaboration

Phys.Rev.Lett.82:3576-3580,1999.

111) [Measurement of b anti-b rapidity correlations in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.D61:032001,2000.

112) [Search for R-parity violating supersymmetry using like sign dielectrons in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.83:2133-2138,1999. [hep-ex/9908063]

113) [Events with a rapidity gap between jets in anti-p p collisions at  \$\sqrt{s} = 630\$ -GeV.](#)

By CDF Collaboration

Phys.Rev.Lett.81:5278-5283,1998.

114) [Search for third generation leptoquarks from technicolor models in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.82:3206,1999.

115) [Kinematics of t anti-t events at CDF.](#)

By CDF Collaboration

Phys.Rev.D59:092001,1999.

116) [Search for a technicolor omega\(T\) particle in events with a photon and a b quark jet at CDF.](#)

By CDF Collaboration

Phys.Rev.Lett.83:3124-3129,1999. [hep-ex/9810031]

117) [Measurement of the top quark mass with the collider detector at Fermilab.](#)

By CDF Collaboration

Phys.Rev.Lett.82:271-276,1999, Erratum-ibid.82:2808-2809,1999. [hep-ex/9810029]

118) [Measurement of  \$Z^0\$  and Drell-Yan production cross-section using dimuons in anti-p p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.D59:052002,1999.

119) [Search for new particles decaying to b anti-b in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.82:2038-2043,1999. [hep-ex/9809022]

120) [Search for the decays  \$B^0\(S\)\$ ,  \$B^0\(d\) \rightarrow e^+ \mu^-\$  and Pati-Salam leptoquarks.](#)

By CDF Collaboration

Phys.Rev.Lett.81:5742-5747,1998.

121) [A Measurement of the lepton charge asymmetry in W boson decays produced in p anti-p collisions.](#)

By CDF Collaboration

Phys.Rev.Lett.81:5754-5759,1998. [hep-ex/9809001]

122) [Search for Higgs bosons produced in association with a vector boson in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.81:5748-5753,1998.

123) [Measurement of the  \$B^0\(s\)\$  meson lifetime using semileptonic decays.](#)

By CDF Collaboration

Phys.Rev.D59:032004,1999. [hep-ex/9808003]



- 124) [Search for second generation leptoquarks in the dimuon plus dijet channel of p anti-p collisions at  \$S^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.81:4806-4811,1998. [hep-ex/9810024]
- 125) [Searches for new physics in diphoton events in p anti-p collisions at  \$s^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D59:092002,1999. [hep-ex/9806034]
- 126) [Measurement of the  \$B\_0\(d\)\$  - anti- \$B\_0\(d\)\$  flavor oscillation frequency and study of same side flavor tagging of B mesons in p anti-p collisions.](#)  
By CDF Collaboration  
Phys.Rev.D59:032001,1999. [hep-ex/9806026]
- 127) [Measurement of the CP violation parameter  \$\sin\(2\text{Beta}\)\$  in  \$B\_0\(d\)/\text{anti-}B\_0\(d\) \rightarrow J/\psi K\_0\(S\)\$  decays.](#)  
By CDF Collaboration  
Phys.Rev.Lett.81:5513-5518,1998. [hep-ex/9806025]
- 128) [Improved measurement of the B- and anti- \$B\_0\$  meson lifetimes using semileptonic decays.](#)  
By CDF Collaboration  
Phys.Rev.D58:092002,1998. [hep-ex/9806018]
- 129) [Observation of the B\(c\) meson in p anti-p collisions at  \$S^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.81:2432-2437,1998. [hep-ex/9805034]
- 130) [Search for longlived parents of  \$Z\_0\$  bosons in p anti-p collisions at  \$S^{1/2} = 1.8\text{-TeV}\$ .](#)  
By The CDF Collaboration  
Phys.Rev.D58:051102,1998. [hep-ex/9805017]
- 131) [Measurement of the  \$\sigma\(W + \geq 1 \text{ jet}\) / \sigma\(W\)\$  cross-section ratio from anti-p p collisions at  \$s^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaborations  
Phys.Rev.Lett.81:1367-1372,1998.
- 132) [Observation of B\(c\) mesons in p anti-p collisions at  \$S^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D58:112004,1998. [hep-ex/9804014]
- 133) [Search for the rare decay  \$W^{+-} \rightarrow D^{+\(-\)}\(s\) \gamma\$  in p anti-p collisions at  \$S^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D58:091101,1998.
- 134) [Search for the rare decay  \$W^{+-} \rightarrow \pi^{+\(-\)} + \gamma\$  in proton - anti-proton collisions at  \$S^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D58:031101,1998.
- 135) [Observation of  \$B^+ \rightarrow \psi\(2S\) K^+\$  and  \$B\_0 \rightarrow \psi\(2S\) K^{\*0}\(892\) \rightarrow 0\$  decays and measurements of B meson branching fractions into  \$J/\psi\$  and  \$\psi\(2S\)\$  final states.](#)  
By CDF Collaboration  
Phys.Rev.D58:072001,1998. [hep-ex/9803013]
- 136) [Search for chargino - neutralino associated production at the Fermilab Tevatron collider.](#)  
By CDF collaboration  
Phys.Rev.Lett.80:5275-5280,1998. [hep-ex/9803015]
- 137) [Searches for new physics in diphoton events in p anti-p collisions at  \$s^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.Lett.81:1791-1796,1998. [hep-ex/9801019]
- 138) [Search for the decays  \$B\(d\)0 \rightarrow \mu^+ \mu^-\$  and  \$B\(s\)0 \rightarrow \mu^+ \mu^-\$  in p anti-p collisions at  \$s^{1/2} = 1.8\text{-TeV}\$ .](#)  
By CDF Collaboration  
Phys.Rev.D57:3811-3816,1998.
- 139) [Observation of hadronic W decays in t anti-t events with the collider detector at Fermilab.](#)  
By CDF Collaboration  
Phys.Rev.Lett.80:5720-5725,1998. [hep-ex/9711004]

140) [Measurement of B hadron lifetimes using J / psi final states at CDF.](#)

By CDF Collaboration

Phys.Rev.D57:5382-5401,1998.

141) [Measurement of the top quark mass and t anti-t production cross-section from dilepton events at the collider detector at Fermilab.](#)

By CDF Collaboration

Phys.Rev.Lett.80:2779-2784,1998. [hep-ex/9802017]

142) [Measurement of the top quark mass.](#)

By CDF Collaboration

Phys.Rev.Lett.80:2767-2772,1998. [hep-ex/9801014]

143) [Measurement of the t anti-t production cross-section in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.80:2773-2778,1998. [hep-ex/9710008]

144) [Measurement of the B0 anti-B0 oscillation frequency in p anti-p collisions using pi B meson charge flavor correlations at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.80:2057-2062,1998. [hep-ex/9712004]

145) [Search for flavor changing neutral current decays of the top quark in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.80:2525-2530,1998.

146) [Properties of jets in W boson events from 1.8-TeV anti-p p collisions.](#)

By CDF Collaboration

Phys.Rev.Lett.79:4760-4765,1997. [hep-ex/9709016]

147) [Measurement of the differential cross-section for events with large total transverse energy in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.80:3461-3466,1998.

148) [Dijet production by color - singlet exchange at the Fermilab Tevatron.](#)

By CDF Collaboration

Phys.Rev.Lett.80:1156-1161,1998.

149) [Search for first generation leptoquark pair production in p anti-p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.79:4327-4332,1997. [hep-ex/9708017]

150) [Search for new particles decaying into b anti-b produced in association with W bosons decaying into e neutrino or mu neutrino at the Tevatron.](#)

By CDF Collaboration

Phys.Rev.Lett.79:3819-3824,1997.

151) [Limits on quark - lepton compositeness scales from dileptons produced in 1.8-TeV p anti-p collisions.](#)

By CDF Collaboration

Phys.Rev.Lett.79:2198-2203,1997.

152) [Search for new gauge bosons decaying into dileptons in anti-p p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.Lett.79:2192-2197,1997.

153) [Properties of photon plus two jet events in anti-p p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration

Phys.Rev.D57:67-77,1998.

154) [Properties of six jet events with large six jet mass at the Fermilab proton - anti-proton collider.](#)

By CDF Collaboration

Phys.Rev.D56:2532-2543,1997.

155) [Double parton scattering in anti-p p collisions at  \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration  
Phys.Rev.D56:3811-3832,1997.

156) [The mu tau and e tau decays of top quark pairs produced in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)

By CDF Collaboration  
Phys.Rev.Lett.79:3585-3590,1997. [hep-ex/9704007]

157) [Measurement of double parton scattering in anti-p p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)

By CDF Collaboration  
Phys.Rev.Lett.79:584-589,1997.

158) [Search for charged Higgs decays of the top quark using hadronic decays of the tau lepton.](#)

By CDF Collaboration  
Phys.Rev.Lett.79:357-362,1997. [hep-ex/9704003]

159) [First observation of the all hadronic decay of t anti-t pairs.](#)

By CDF Collaboration  
Phys.Rev.Lett.79:1992-1997,1997.

160) [Measurement of diffractive dijet production at the Tevatron.](#)

By CDF Collaboration  
Phys.Rev.Lett.79:2636-2641,1997.

161) [The CDF-II detector: Technical design report.](#)

By CDF-II Collaboration  
FERMILAB-PUB-96-390-E (Nov 1996) 234p.

162) [Production of J / psi mesons from chi\(c\) meson decays in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)

By CDF Collaboration  
Phys.Rev.Lett.79:578-583,1997.

163) [Search for gluinos and squarks at the Fermilab Tevatron Collider.](#)

By CDF Collaboration  
Phys.Rev.D56:1357-1362,1997.

164) [J / psi and psi \(2S\) production in p anti-p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)

By CDF Collaboration  
Phys.Rev.Lett.79:572-577,1997.

165) [Observation of diffractive W boson production at the Tevatron.](#)

By CDF Collaboration  
Phys.Rev.Lett.78:2698-2703,1997. [hep-ex/9703010]

166) [Search for new particles decaying to dijets at CDF.](#)

By CDF Collaboration  
Phys.Rev.D55:5263-5268,1997. [hep-ex/9702004]

167) [Search for third generation leptoquarks in anti-p p collisions at  \$s^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)

By CDF Collaboration  
Phys.Rev.Lett.78:2906-2911,1997.

168) [Measurement of dijet angular distributions at CDF.](#)

By CDF Collaboration  
Phys.Rev.Lett.77:5336-5341,1996, Erratum-ibid.78:4307,1997. [hep-ex/9609011]

169) [Observation of W+ W- production in anti-p p collisions at  \$S^{\*\*}\(1/2\) = 1.8\text{-TeV}\$ .](#)

By CDF Collaboration  
Phys.Rev.Lett.78:4536-4540,1997.

170) [Measurement of the branching fraction B \(B\(u\)+ ---> J / psi pi+\) and search for B\(c\)+ ---> J / psi pi+.](#)

By CDF Collaboration  
Phys.Rev.Lett.77:5176-5181,1996.

171) [Observation of Lambda\(b\)0 ---> J / psi Lambda at the Fermilab Proton - Anti-proton Collider.](#)

By CDF Collaboration  
Phys.Rev.D55:1142-1152,1997.

172) [Measurement of b anti-b production correlations, B0 anti-B0 mixing, and a limit on epsilon\(B\) in p anti-p collisions at S\\*\\*\(1/2\) = 1.8-TeV.](#)

By CDF Collaboration  
Phys.Rev.D55:2546-2558,1997.

173) [Charmed mesons fragmentation functions.](#)

By M. Cacciari, M. Greco, S. Rolli, A. Tanzini.  
Phys.Rev.D55:2736-2740,1997. [hep-ph/9608213]

174) [Fragmentation functions approach in pQCD fragmentation phenomena.](#)

By Simona Rolli.  
In \*Les Arcs 1996, QCD and high energy hadronic interactions\* 249-25. [hep-ph/9607480]

175) [Top mass measurement at CDF.](#)

By Simona Rolli.  
In \*Padua 1996, Hadron collider physics\* 449- 46.

176) [Ratios of bottom meson branching fractions involving j / psi mesons and determination of b quark fragmentation fractions.](#)

By CDF Collaboration  
Phys.Rev.D54:6596-6609,1996. [hep-ex/9607003]

177) [Further properties of high mass multi - jet events at the Fermilab proton - anti-proton collider.](#)

By CDF Collaboration  
Phys.Rev.D54:4221-4233,1996. [hep-ex/9605004]

178) [Measurement of the lifetime of the B\(s\)0 meson using the exclusive decay mode B\(s\)0 ---> J / psi phi.](#)

By CDF Collaboration  
Phys.Rev.Lett.77:1945-1949,1996.

179) [Measurement of Lambda\(b\)0 lifetime using Lambda\(b\)0 ---> Lambda\(c\)+ lepton- anti-neutrino.](#)

By CDF Collaboration  
Phys.Rev.Lett.77:1439-1443,1996.

180) [Properties of jets in Z boson events from 1.8-TeV anti-p p collisions.](#)

By CDF Collaboration  
Phys.Rev.Lett.77:448-453,1996. [hep-ex/9603003]

181) [Search for flavor changing neutral current B meson decays in p anti-p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.76:4675-4680,1996.

182) [Measurement of the B- and anti-B0 meson lifetimes using semileptonic decays.](#)

By CDF Collaboration  
Phys.Rev.Lett.76:4462-4467,1996.

183) [Search for chargino - neutralino production in p anti-p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.76:4307-4311,1996. [hep-ex/9603001]

184) [Inclusive jet cross-section in anti-p p collisions at s\\*\\*\(1/2\) = 1.8-TeV.](#)

By CDF Collaboration  
Phys.Rev.Lett.77:438-443,1996. [hep-ex/9601008]

185) [Search for charged Higgs decays of the top quark using hadronic tau decays.](#)

By CDF Collaboration  
Phys.Rev.D54:735-742,1996. [hep-ex/9601003]

186) [Kinematical probes of neutrino mass.](#)

By N1 working Group  
In \*Snowmass 1994, Particle and nuclear astrophysics and cosmology in the next millennium\* 195-21.

187) [Light mesons production at the tevatron to next-to-leading order.](#)

By Mario Greco & Simona Rolli.

Phys.Rev.D52:3853-3861,1995. [hep-ph/9409356]

188) [Inclusive particle photoproduction to next-to-leading order.](#)

By M. Greco, S. Rolli, A. Vicini.

Z.Phys.C65:277-284,1995. [hep-ph/9404228]

189) [Next-to-leading order eta production at hadron colliders.](#)

By M. Greco & S. Rolli.

Z.Phys.C60:169-174,1993. [hep-ph/9304311]

190) [Next-to-leading order determination of pion fragmentation functions.](#)

By P. Chiappetta, M. Greco, J.P. Guillet, S. Rolli, M. Werlen.

Nucl.Phys.B412:3-38,1994. [hep-ph/9301254]

---

Link to the [main SPIRES-HEP search page](#)

[Search HEP](#) | [Other Databases](#) | [Conferences](#) | [Institutions](#) | [Experiments](#) | [Help](#) | [Comments](#)  
[SLAC](#) | [SLAC Library](#) | [Stanford University](#)



Questions and Comments to [library@slac.stanford.edu](mailto:library@slac.stanford.edu)

Location: SPIRES HEP Results Display